

Theory of Neutrino Oscillations¹

Carlo Giunti

INFN, Sezione di Torino, and Dipartimento di Fisica Teorica,
Università di Torino, Via P. Giuria 1, I-10125 Torino, Italy

Abstract

We review critically the main assumptions on which the standard theory of neutrino oscillations is based. We show that all assumptions are realistic, except the so-called “equal momentum assumption”, which however is irrelevant. We briefly review the covariant plane-wave derivation of neutrino oscillations and a quantum field theoretical wave packet model of neutrino oscillations. We show that both approaches lead to the standard expression for the oscillation phase. The wave packet model allows also to describe the coherence of the oscillations and the localization of the production and detection processes.

1 Introduction

The possibility of neutrino oscillations was discovered by Bruno Pontecorvo in the late 50's following an analogy with kaon oscillations [1,2]. Since at that time only one *active* neutrino was known, Pontecorvo invented the concept of a *sterile* neutrino [3], which is a neutral fermion which does not take part to weak interactions. The muon neutrino was discovered in 1962 in the Brookhaven experiment of Lederman, Schwartz, Steinberger *et al.* [4], which followed a proposal made by Pontecorvo in 1959 [5]. Then, it became clear that oscillations between different active neutrino flavors are possible if neutrinos are massive and mixed particles. Indeed, in 1967 [3] Pontecorvo predicted the solar neutrino problem as a possible result of $\nu_e \rightarrow \nu_\mu$ (or $\nu_e \rightarrow \nu_{\text{sterile}}$) transitions before the first measurement of the solar electron neutrino flux in the Homestake experiment [6], and in 1969 Gribov and Pontecorvo discussed in detail the possibility of solar neutrino oscillations due to neutrino mixing [7].

However, in these and other papers written before 1976 the probability of neutrino oscillations was not calculated in a rigorous way, but simply estimated on the basis of the analogy with kaon oscillations. As a result, the phase of the oscillations was correct within a factor of two.

The standard theory of neutrino oscillations was developed in 1976 by Eliezer and Swift [8], Fritzsche and Minkowski [9], Bilenky and Pontecorvo [10] (see the beautiful

¹Talk presented at the 11th Lomonosov Conference on Elementary Particle Physics, 21–27 August 2003, Moscow State University, Moscow, Russia.

review in Ref. [11]) on the basis of the following four assumptions that will be discussed critically in this report:

(A1) Neutrinos are ultrarelativistic particles.

(A2) Neutrinos produced or detected in CC weak interaction processes are described by the flavor states

$$|\nu_\alpha\rangle = \sum_k U_{\alpha k}^* |\nu_k\rangle, \quad (1.1)$$

where U is the unitary mixing matrix, $\alpha = e, \mu, \tau$, and $|\nu_k\rangle$ is the state of a neutrino with mass m_k .

(A3) The propagation time is equal to the distance L traveled by the neutrino between production and detection.

(A4) The massive neutrino states $|\nu_k\rangle$ in Eq. (1.1) have the same momentum, $p_k = p \simeq E$ (“equal momentum assumption”), and different energies, $E_k = \sqrt{p^2 + m_k^2} \simeq E + m_k^2/2E$, where E is the neutrino energy neglecting mass effects and the approximations are valid for ultrarelativistic neutrinos.

In Section 2 we briefly review the main points of the standard theory of neutrino oscillations. In Sections 3–6 we discuss critically the four assumptions listed above. In Section 7 we review the covariant derivation of the neutrino oscillation probability in the plane wave approach [12–15]. In Section 8 we review the quantum field theoretical wave packet model presented in Ref. [16]. Finally, in Section 9 we present our conclusions.

2 Standard theory of neutrino oscillations

In the plane wave approximation the states $|\nu_k\rangle$ of massive neutrinos are eigenstates of the free Hamiltonian with definite energy eigenvalues E_k . Therefore, their time evolution is given by the Schrödinger equation, whose solution is

$$|\nu_k(t)\rangle = e^{-iE_k t} |\nu_k\rangle. \quad (2.1)$$

Using assumption (A2), from Eq. (1.1) the time evolution of the flavor states is given by

$$|\nu_\alpha(t)\rangle = \sum_{\beta=e,\mu,\tau} \left(\sum_k U_{\alpha k}^* e^{-iE_k t} U_{\beta k} \right) |\nu_\beta\rangle, \quad (2.2)$$

which, for $t > 0$, is a superposition of different flavors if the mixing matrix is non-diagonal. The coefficient of the flavor state $|\nu_\beta\rangle$ is the amplitude of $\nu_\alpha \rightarrow \nu_\beta$ transitions, whose squared absolute value gives the probability

$$P_{\nu_\alpha \rightarrow \nu_\beta}(t) = |\langle \nu_\beta | \nu_\alpha(t) \rangle|^2 = \left| \sum_k U_{\alpha k}^* e^{-iE_k t} U_{\beta k} \right|^2. \quad (2.3)$$

Using the equal-momentum assumption (A4), the energy of the k^{th} massive neutrino component is given by $E_k = \sqrt{p^2 + m_k^2}$, which can be approximated to $E_k \simeq p + m_k^2/2p$

in the case of ultrarelativistic neutrinos following from assumption (A1). Moreover, assumption (A3) allows to replace the usually unknown propagation time t with the usually known distance L traveled by the neutrino between production and detection. The final result for the oscillation probability can be written as

$$P_{\nu_\alpha \rightarrow \nu_\beta}(L) = \sum_k |U_{\alpha k}|^2 |U_{\beta k}|^2 + 2\text{Re} \sum_{k>j} U_{\alpha k}^* U_{\beta k} U_{\alpha j} U_{\beta j}^* \exp\left(-i \frac{\Delta m_{kj}^2 L}{2E}\right), \quad (2.4)$$

where $\Delta m_{kj}^2 = m_k^2 - m_j^2$ and $E = p$ is the neutrino energy neglecting mass contributions. In Eq. (2.4) we have separated the expression for the flavor transition probability into a constant term and a term which oscillates as a function of the distance L . The oscillating term is the most interesting one from a quantum mechanical point of view, because it is due to the interference between the different massive neutrino components, whose existence requires coherent production and detection. On the other hand, the constant term is experimentally very important, because it gives the average probability of flavor transitions, which is the measured one when the oscillating term is not present because of lack of coherence or when the oscillating term is not measurable because it is washed out by the average over the energy resolution of the detector or the distance uncertainty.

Let us now examine critically one by one the four assumptions (A1)–(A4) that led to the result (2.4).

3 Assumption (A1): ultrarelativistic neutrinos

The assumption (A1) is correct, because neutrino masses are smaller than about one eV (see Refs. [17, 18]) and only neutrinos with energy larger than about 100 keV can be detected.

Indeed, neutrinos are detected in:

1. Charged-current or neutral-current weak processes which have an energy threshold larger than some fraction of MeV. This is due to the fact that in a scattering process $\nu + A \rightarrow \sum_X X$ with A at rest, the squared center-of-mass energy $s = 2E_\nu m_A + m_A^2$ (neglecting the neutrino mass) must be bigger than $(\sum_X m_X)^2$, leading to $E_\nu^{\text{th}} = \frac{(\sum_X m_X)^2}{2m_A} - \frac{m_A}{2}$. For example:
 - $E_\nu^{\text{th}} \simeq 0.233 \text{ MeV}$ for $\nu_e + {}^{71}\text{Ga} \rightarrow {}^{71}\text{Ge} + e^-$ in gallium solar neutrino experiments (see Ref. [18]).
 - $E_\nu^{\text{th}} \simeq 0.81 \text{ MeV}$ for $\nu_e + {}^{37}\text{Cl} \rightarrow {}^{37}\text{Ar} + e^-$ in the Homestake [6] solar neutrino experiment.
 - $E_\nu^{\text{th}} \simeq 1.8 \text{ MeV}$ for $\bar{\nu}_e + p \rightarrow n + e^+$ in reactor neutrino experiments (see Ref. [18]).
 - $E_\nu^{\text{th}} \simeq 2.2 \text{ MeV}$ in the neutral-current process $\nu + d \rightarrow p + n + \nu$ used in the SNO experiment to detect active solar neutrinos [19].
2. The elastic scattering process $\nu + e^- \rightarrow \nu + e^-$, whose cross section is proportional to the neutrino energy ($\sigma(E_\nu) \sim \sigma_0 E_\nu / m_e$, with $\sigma_0 \sim 10^{-44} \text{ cm}^2$). An energy threshold of some MeV's is needed in order to have a signal above the background. For example, $E_\nu^{\text{th}} \simeq 5 \text{ MeV}$ in the Super-Kamiokande [20] solar neutrino experiment.

As we will see, the ultrarelativistic character of neutrinos implies the correctness of the assumptions (A2) and (A3) and the irrelevance of the assumption (A4), which is not realistic.

4 Assumption (A2): flavor states

In Ref. [21] it has been shown that the assumption (A2) is not exact, because the amplitude of production and detection of the massive neutrino ν_k is not simply given by $U_{\alpha k}^*$ (see also Refs. [14, 16]). However, in the ultrarelativistic approximation the characteristics of the production and detection processes that depend on the neutrino mass can be neglected, leading to a correct approximate description of flavor neutrinos through the states (1.1).

5 Assumption (A3): $t = L$

The assumption (A3) follows from the ultrarelativistic approximation, because ultrarelativistic particles propagate almost at the velocity of light. However, in the standard theory of neutrino oscillations massive neutrinos are treated as plane waves, which are limitless in space and time. In order to justify the assumption (A3) it is necessary to treat massive neutrinos as wave packets [22], which are localized on the production process at the production time and propagate between the production and detection processes at a velocity close to the velocity of light. Such a wave packet treatment [16, 22–24] yields the standard formula for the oscillation length. In addition, the different group velocities of different massive neutrinos imply the existence of a coherence length for the oscillations, beyond which the wave packets of different massive neutrinos do not jointly overlap with the detection process [25, 26].

The wave packet treatment of neutrino oscillations is also necessary for a correct description of the momentum and energy uncertainties necessary for the coherent production and detection of different massive neutrinos [24, 27, 28], whose interference generates the oscillations.

The physical reason why the substitution $t = L$ is correct can be understood by noting that, if the massive neutrinos are ultrarelativistic and contribute coherently to the detection process, their wave packets overlap with the detection process for an interval of time $[t - \Delta t, t + \Delta t]$, with

$$t = \frac{L}{\bar{v}} \simeq L \left(1 + \frac{\overline{m^2}}{2E^2} \right), \quad \Delta t \sim \sigma_x, \quad (5.1)$$

where \bar{v} is the average group velocity, $\overline{m^2}$ is the average of the squared neutrino masses, σ_x is given by the spatial uncertainties of the production and detection processes summed in quadrature [23] (the spatial uncertainty of the production process determines the size of the massive neutrino wave packets). The correction $L\overline{m^2}/2E^2$ to $t = L$ in Eq. (5.1) can be neglected, because it gives corrections to the oscillation phases which are of higher order in the very small ratios m_k^2/E^2 . The corrections due to $\Delta t \sim \sigma_x$ are also negligible, because in all realistic experiments σ_x is much smaller than the oscillation length

$L_{kj}^{\text{osc}} = 4\pi E/\Delta m_{kj}^2$, otherwise oscillations could not be observed [22, 24, 27, 28]. One can summarize these arguments by saying that the substitution $t = L$ is correct because the phase of the oscillations is practically constant over the interval of time in which the massive neutrino wave packets overlap with the detection process and it is given by

$$\phi_{kj}(L) = \frac{\Delta m_{kj}^2 L}{2E} = 2\pi \frac{L}{L_{kj}^{\text{osc}}}, \quad (5.2)$$

plus negligible corrections of higher order in the neutrino masses.

6 Assumption (A4): equal momentum

Let us discuss now the assumption (A4), which has been shown to be unrealistic in Refs. [24, 29] on the basis of simple relativistic arguments. Indeed, the relativistic transformation of energy and momentum implies that the equal momentum assumption cannot hold concurrently in different inertial systems. On the other hand, the probability of flavor neutrino oscillations is independent from the inertial system adopted for its measurement, because the neutrino flavor is measured by observing charged leptons whose character is Lorentz invariant (*e.g.* an electron is seen as an electron in any system of reference). Therefore, the probability of neutrino oscillations is Lorentz invariant [13, 30] and must be derived in a covariant way. In fact, the oscillation probability has been derived without special assumptions about the energies and momenta of the different massive neutrino components both in the plane wave approach [12–15] and in the wave packet treatment [22–24, 29].

7 Covariant derivation of neutrino oscillations

Let us briefly describe the covariant derivation of the neutrino oscillation probability in the plane wave approach, in which the massive neutrino states in Eq. (1.1) evolve in space and time as plane waves:

$$|\nu_k(x, t)\rangle = e^{-iE_k t + ip_k x} |\nu_k\rangle. \quad (7.1)$$

Substituting Eq. (7.1) in Eq. (1.1) and expressing the $|\nu_k\rangle$ on the right-hand side in terms of flavor states ($|\nu_k\rangle = \sum_{\beta=e,\mu,\tau} U_{\beta k}^* |\nu_\beta\rangle$), we obtain

$$|\nu_\alpha(x, t)\rangle = \sum_{\beta=e,\mu,\tau} \left(\sum_k U_{\alpha k} e^{-iE_k t + ip_k x} U_{\beta k}^* \right) |\nu_\beta\rangle, \quad (7.2)$$

which shows that at a distance x and after a time t from the production of a neutrino with flavor α , the neutrino is a superposition of different flavors (if the mixing matrix is not diagonal). The probability of flavor transitions in space and time is given by

$$P_{\nu_\alpha \rightarrow \nu_\beta}(x, t) = |\langle \nu_\beta | \nu_\alpha(x, t) \rangle|^2 = \left| \sum_k U_{\alpha k} e^{-iE_k t + ip_k x} U_{\beta k}^* \right|^2, \quad (7.3)$$

which is manifestly Lorentz invariant.

Considering ultrarelativistic neutrinos, we apply now the assumption (A3), $t = x = L$, where L is the distance traveled by the neutrino between production and detection. The phase in Eq. (7.3) becomes

$$E_k t - p_k x = (E_k - p_k) L = \frac{E_k^2 - p_k^2}{E_k + p_k} L = \frac{m_k^2}{E_k + p_k} L \simeq \frac{m_k^2}{2E} L. \quad (7.4)$$

It is important to notice that Eq. (7.4) shows that the phases of massive neutrinos relevant for the oscillations are independent from any assumption on the energies and momenta of different massive neutrinos, as long as the relativistic dispersion relation $E_k^2 = p_k^2 + m_k^2$ is satisfied. This is why the standard derivation of the neutrino oscillation probability gives the correct result, in spite of the unrealistic equal momentum assumption (A4).

Using the phase in Eq. (7.4), the oscillation probability as a function of the distance L has the standard expression in Eq. (2.4). Let us notice that the expression (2.4) is still Lorentz invariant, as shown in Ref. [30], because L is not the instantaneous source-detector distance but the distance traveled by the neutrino between production and detection.

8 Wave packet model

Several wave packet models of neutrino oscillations have been devised, with similar results, in the framework of Quantum Mechanics [22–24, 26, 29, 31] and Quantum Field Theory [16, 32–35] (see Ref. [28] for a comprehensive review). Here we briefly review the main points of the quantum field theoretical wave packet model presented in Ref. [16], which is based on the assumption in Quantum Field Theory that free particles are described by wave packets constructed as appropriate superpositions of states in the momentum Fock space of the corresponding free field.

The wave packet describing a neutrino created with flavor α in the process

$$P_I \rightarrow P_F + \ell_\alpha^+ + \nu_\alpha \quad (8.1)$$

is given by

$$|\nu_\alpha\rangle \propto \langle P_F, \ell_\alpha^+ | -i \int d^4x \mathcal{H}_I(x) |P_I\rangle, \quad (8.2)$$

where we have considered the first order perturbative contribution of the effective weak interaction hamiltonian $\mathcal{H}_I(x)$. The states $|P_I\rangle$, $|P_F\rangle$, $|\ell_\alpha^+\rangle$ that describe the particles taking part to the localized production process have the wave packet form

$$|\chi\rangle = \int d^3p \psi_\chi(\vec{p}; \vec{p}_\chi, \sigma_{p\chi}) |\chi(\vec{p}, h_\chi)\rangle \quad (\chi = P_I, P_F, \ell_\alpha^+), \quad (8.3)$$

where \vec{p}_χ is the average momentum, $\sigma_{p\chi}$ is the momentum uncertainty, determined by the interactions with the surrounding medium, and h_χ is the helicity. Approximating the momentum distributions $\psi_\chi(\vec{p}; \vec{p}_\chi, \sigma_{p\chi})$ with gaussian functions, the integrals in the expression (8.2) for the neutrino state can be calculated analytically, leading to

$$|\nu_\alpha\rangle = N_\alpha \sum_k U_{\alpha k}^* \int d^3p e^{-S_k^P(\vec{p})} \sum_h \mathcal{A}_k^P(\vec{p}, h) |\nu_k(\vec{p}, h)\rangle, \quad (8.4)$$

where N_α is a normalization factor, $\mathcal{A}_k^P(\vec{p}, h)$ is the amplitude of production of a neutrino with mass m_k , momentum \vec{p} and helicity h , and the function $e^{-S_k^P(\vec{p})}$ enforces energy-momentum conservation within the momentum uncertainty due to the momentum distributions of P_I, P_F, ℓ_α^+ . The spatial width σ_{xP} of the neutrino wave packet is related to its momentum uncertainty

$$\sigma_{pP}^2 = \sigma_{pP_I}^2 + \sigma_{pP_F}^2 + \sigma_{p\ell_\alpha^+}^2 \quad (8.5)$$

by the minimal Heisenberg uncertainty relation $\sigma_{xP}\sigma_{pP} = 1/2$.

Let us consider the detection of a neutrino with flavor β through the charged-current weak process

$$\nu_\beta + D_I \rightarrow D_F + \ell_\beta^-, \quad (8.6)$$

at a space-time distance (\vec{L}, T) from the production process, where the neutrino created with flavor α is described by a state obtained by acting on $|\nu_\alpha\rangle$ in Eq. (8.4) with the space-time translation operator $\exp(-i\widehat{E}T + i\widehat{P}\cdot\vec{L})$, where \widehat{E} and \widehat{P} are the energy and momentum operators, respectively:

$$|\nu_\alpha(\vec{L}, T)\rangle = N_\alpha \sum_k U_{\alpha k}^* \int d^3p e^{-iE_{\nu_k}(\vec{p})T + i\vec{p}\cdot\vec{L}} e^{-S_k^P(\vec{p})} \sum_h \mathcal{A}_k^P(\vec{p}, h) |\nu_k(\vec{p}, h)\rangle. \quad (8.7)$$

The detection amplitude is given by

$$\mathcal{A}_{\alpha\beta}(\vec{L}, T) = \langle D_F, \ell_\beta^- | -i \int d^4x \mathcal{H}_I(x) | D_I, \nu_\alpha(\vec{L}, T) \rangle. \quad (8.8)$$

The transition probability as a function of the distance \vec{L} is given by the average over the unmeasured time T of $|\mathcal{A}_{\alpha\beta}(\vec{L}, T)|^2$. In the realistic case of ultrarelativistic neutrinos the final result for the flavor transition probability is

$$P_{\alpha\beta}(L) = \sum_k |U_{\alpha k}|^2 |U_{\beta k}|^2 + 2\text{Re} \sum_{k>j} U_{\alpha k}^* U_{\beta k} U_{\alpha j} U_{\beta j}^* \exp \left[-2\pi i \frac{L}{L_{kj}^{\text{osc}}} - \left(\frac{L}{L_{kj}^{\text{coh}}} \right)^2 - 2\pi^2 \kappa \left(\frac{\sigma_x}{L_{kj}^{\text{osc}}} \right)^2 \right], \quad (8.9)$$

where $L_{kj}^{\text{osc}} = 4\pi E / \Delta m_{kj}^2$ are the standard oscillation lengths and $L_{kj}^{\text{coh}} = 4\sqrt{2}\omega E^2 \sigma_x / |\Delta m_{kj}^2|$ are the coherence lengths. The quantities κ and ω , which are usually of order one, depend on the production and detection processes [16]. The total spatial coherence width σ_x is given by

$$\sigma_x^2 = \sigma_{xP}^2 + \sigma_{xD}^2. \quad (8.10)$$

The form of the flavor transition probability in Eq. (8.9) is consistent with the results obtained with other wave packet models in the framework of Quantum Mechanics [22–24, 26, 29, 31] and Quantum Field Theory [32–35]. One can see that the standard value of the oscillation phase is confirmed, in agreement with the discussion in the previous Sections. In addition, the wave packet treatment produced a coherence term and a localization term.

The coherence term $\exp[-(L/L_{kj}^{\text{coh}})^2]$ suppresses the oscillations due to Δm_{kj}^2 when $L \gtrsim L_{kj}^{\text{coh}}$, because the wave packets of the massive neutrino components ν_k and ν_j have separated so much that they cannot be absorbed coherently in the detection process.

The localization term $\exp[-2\pi^2\kappa(\sigma_x/L_{kj}^{\text{osc}})^2]$ suppresses the oscillations due to Δm_{kj}^2 if $\sigma_x \gtrsim L_{kj}^{\text{osc}}$. This means that in order to measure the interference of the massive neutrino components ν_k and ν_j the production and detection processes must be localized in space-time regions much smaller than the oscillation length L_{kj}^{osc} . In practice this requirement is easily satisfied by all neutrino oscillation experiments, because the space-time coherence regions of the production and detection processes are usually microscopic, whereas the oscillation length is usually macroscopic.

The localization term is important for the distinction of neutrino oscillation experiments from experiments on the measurement of neutrino masses. As first shown by Kayser in Ref. [27], neutrino oscillations are suppressed in experiments able to measure the value of a neutrino mass, because the measurement of a neutrino mass implies that only the corresponding massive neutrino is produced or detected.

Kayser's [27] argument goes as follows. Since a neutrino mass is measured from energy-momentum conservation in a process in which a neutrino is produced or detected, from the energy-momentum dispersion relation $E_k^2 = p_k^2 + m_k^2$ the uncertainty of the mass determination is

$$\delta m_k^2 = \sqrt{(2E_k\delta E_k)^2 + (2p_k\delta p_k)^2} \simeq 2\sqrt{2}E\sigma_p, \quad (8.11)$$

where the approximation holds for realistic ultrarelativistic neutrinos and $\sigma_p = 1/2\sigma_x$ is the momentum uncertainty. If $\delta m_k^2 < |\Delta m_{kj}^2|$, the mass of ν_k is measured with an accuracy better than the difference Δm_{kj}^2 . In this case the neutrino ν_j is not produced or detected and the interference of ν_k and ν_j is not observed. The localization term in the oscillation probability (8.9) automatically implements Kayser's mechanism, because $\sigma_x/L_{kj}^{\text{osc}}$ can be written as $\Delta m_{kj}^2/4\sqrt{2}E\sigma_p$. If $\delta m_k^2 < |\Delta m_{kj}^2|$, the localization term in Eq. (8.9) suppresses² the interference of ν_k and ν_j .

9 Conclusions

In conclusion, we have shown that the probability of neutrino oscillations can be derived in a covariant way in the plane wave approach starting from realistic assumptions. We have also presented a derivation of neutrino oscillations in a quantum field theoretical wave packet approach. In both cases we obtained the standard expression for the oscillation phase. The wave packet approach allows also to describe the coherence of the oscillations and the localization of the production and detection processes.

References

- [1] B. Pontecorvo, Sov. Phys. JETP **6**, 429 (1957).
- [2] B. Pontecorvo, Sov. Phys. JETP **7**, 172 (1958).

²Using the arguments presented by M. Beuthe in Ref. [35], it is possible to show that κ is always of order one.

- [3] B. Pontecorvo, Sov. Phys. JETP **26**, 984 (1968).
- [4] G. Danby *et al.*, Phys. Rev. Lett. **9**, 36 (1962).
- [5] B. Pontecorvo, Sov. Phys. JETP **37**, 1236 (1960).
- [6] Homestake, B. T. Cleveland *et al.*, Astrophys. J. **496**, 505 (1998).
- [7] V. N. Gribov and B. Pontecorvo, Phys. Lett. **B28**, 493 (1969).
- [8] S. Eliezer and A. R. Swift, Nucl. Phys. **B105**, 45 (1976).
- [9] H. Fritzsch and P. Minkowski, Phys. Lett. **B62**, 72 (1976).
- [10] S. M. Bilenky and B. Pontecorvo, Nuovo Cim. Lett. **17**, 569 (1976).
- [11] S. M. Bilenky and B. Pontecorvo, Phys. Rept. **41**, 225 (1978).
- [12] R. G. Winter, Lett. Nuovo Cim. **30**, 101 (1981).
- [13] C. Giunti and C. W. Kim, Found. Phys. Lett. **14**, 213 (2001), [hep-ph/0011074](#).
- [14] S. M. Bilenky and C. Giunti, Int. J. Mod. Phys. **A16**, 3931 (2001), [hep-ph/0102320](#).
- [15] C. Giunti, (2003), [hep-ph/0311241](#), IFAE 2003, Lecce, 23-26 April 2003.
- [16] C. Giunti, JHEP **11**, 017 (2002), [hep-ph/0205014](#).
- [17] S. M. Bilenky, C. Giunti, J. A. Grifols, and E. Masso, Phys. Rept. **379**, 69 (2003), [hep-ph/0211462](#).
- [18] C. Giunti and M. Laveder, (2003), [hep-ph/0310238](#).
- [19] SNO, Q. R. Ahmad *et al.*, Phys. Rev. Lett. **89**, 011301 (2002), [nucl-ex/0204008](#).
- [20] SuperKamiokande, S. Fukuda *et al.*, Phys. Rev. Lett. **86**, 5651 (2001), [hep-ex/0103032](#).
- [21] C. Giunti, C. W. Kim, and U. W. Lee, Phys. Rev. **D45**, 2414 (1992).
- [22] C. Giunti, C. W. Kim, and U. W. Lee, Phys. Rev. **D44**, 3635 (1991).
- [23] C. Giunti and C. W. Kim, Phys. Rev. **D58**, 017301 (1998), [hep-ph/9711363](#).
- [24] C. Giunti, Found. Phys. Lett. **17**, 103 (2004), [hep-ph/0302026](#).
- [25] S. Nussinov, Phys. Lett. **B63**, 201 (1976).
- [26] K. Kiers, S. Nussinov, and N. Weiss, Phys. Rev. **D53**, 537 (1996), [hep-ph/9506271](#).
- [27] B. Kayser, Phys. Rev. **D24**, 110 (1981).
- [28] M. Beuthe, Phys. Rept. **375**, 105 (2003), [hep-ph/0109119](#).
- [29] C. Giunti, Mod. Phys. Lett. **A16**, 2363 (2001), [hep-ph/0104148](#).

- [30] C. Giunti, (2003), [physics/0305122](#).
- [31] K. Kiers and N. Weiss, Phys. Rev. **D57**, 3091 (1998), [hep-ph/9710289](#).
- [32] C. Giunti, C. W. Kim, J. A. Lee, and U. W. Lee, Phys. Rev. **D48**, 4310 (1993), [hep-ph/9305276](#).
- [33] C. Giunti, C. W. Kim, and U. W. Lee, Phys. Lett. **B421**, 237 (1998), [hep-ph/9709494](#).
- [34] C. Y. Cardall, Phys. Rev. **D61**, 073006 (2000), [hep-ph/9909332](#).
- [35] M. Beuthe, Phys. Rev. **D66**, 013003 (2002), [hep-ph/0202068](#).